
BTeV's Staged Detector & Some Physics Reach Comparisons with LHCb

Belarussian State- D .Drobychev,
A. Lobko, A. Lopatrik, R. Zouversky

UC Davis - P. Yager

Univ. of Colorado at Boulder

J. Cumalat, P. Rankin, K. Stenson

Fermi National Lab

J. Appel, E. Barsotti, C. Brown,
J. Butler, H. Cheung, D. Christian,
S. Cihangir, M. Fischler,
I. Gaines, P. Garbincius, L. Garren,
E. Gottschalk, A. Hahn, G. Jackson,
P. Kasper, P. Kasper, R. Kutschke,
S. W. Kwan, P. Lebrun, P. McBride,
J. Slaughter, M. Votava, M. Wang,
J. Yarba

Univ. of Florida at Gainesville
P. Avery

University of Houston –

A. Daniel, K. Lau, M. Ispiryan,
B. W. Mayes, V. Rodriguez,
S. Subramania, G. Xu

Illinois Institute of Technology

R. Burnstein, D. Kaplan,
L. Lederman, H. Rubin, C. White

Univ. of Illinois- M. Haney, D.
Kim, M. Selen, V. Simatis, J. Wiss

Univ. of Insubria in Como-
P. Ratcliffe, M. Rovere

INFN - Frascati- M. Bertani, L.
Benussi, S. Bianco, M. Caponero, D.
Collona, F. Fabri, F. Di Falco, F.
Felli, M. Giardoni, A. La Monaca,
E. Pace, M. Pallota, A. Paolozzi, S.
Tomassini

INFN - Milano – G. Alimonti,
P'Dangelo, M. Dinardo, L. Edera, S.
Erba, D. Lunesu, S. Magni, D.
Menasce, L. Moroni, D. Pedrini, S.
Sala, L. Uplegger

INFN - Pavia - G. Boca, G.
Cossali, G. Liguori, F. Manfredi, M.
Maghisoni, L. Ratti, V. Re, M.
Santini, V. Speviali, P. Torre, G.
Traversi

IHEP Protvino, Russia - A.
Derevschikov, Y. Goncharenko, V.
Khodyrev, V. Kravtsov, A.
Meschanin, V. Mochalov,
D. Morozov, L. Nogach, P.
Semenov K. Shestermanov,
L. Soloviev, A. Uzunian, A. Vasiliev

University of Iowa

C. Newsom, & R. Braunger

University of Minnesota
J. Hietala, Y. Kubota, B. Lang,
R. Poling, A. Smith

Nanjing Univ. (China)-
T. Y. Chen, D. Gao, S. Du,
M. Qi, B. P. Zhang, Z. Xi
Xang, J. W. Zhao

New Mexico State -
V. Papavassiliou

Northwestern Univ. -
J. Rosen

Ohio State University-
K. Honscheid, & H. Kagan

Univ. of Pennsylvania
W. Selove

Univ. of Puerto Rico
A. Lopez, H. Mendez, J.
Ramirez, W. Xiong

**Univ. of Science & Tech.
of China** - G. Datao, L. Hao,
Ge Jin, L. Tiankuan, T. Yang,
& X. Q. Yu

Shandong Univ. (China)-
C. F. Feng, Yu Fu, Mao He,
J. Y. Li, L. Xue, N. Zhang, &
X. Y. Zhang

Southern Methodist –
T. Coan, M. Hosack

Syracuse University-

M. Artuso, C. Boulahouache,
S. Blusk, J. Butt, O.
Dorjkhaidav, J. Haynes, N.
Mena, R. Mountain,
H. Muramatsu, R. Nandakumar,
L. Redjimi, R. Sia,
T. Skwarnicki, S. Stone, J. C.
Wang, K. Zhang

Univ. of Tennessee

T. Handler, R. Mitchell
Vanderbilt University
W. Johns, P. Sheldon,
E. Vaandering, & M. Webster

University of Virginia M.
Arenton, S. Conetti, B. Cox, A.
Ledovskoy, H. Powell, M.
Ronquest, D. Smith, B.
Stephens, Z. Zhe

Wayne State University
G. Bonvicini, D. Cinabro,
A. Schreiner

University of Wisconsin
M. Sheaff

York University - S. Menary

Some Significant Events in B Physics

Year	Item	Theory Prediction	~Value	# B's
1983	τ_b	Too small to be observed $\sim < 0.1$ ps	1 ps	2×10^4
1987	$B^0 - \bar{B}^0$ mixing	Too small to see ($\sim < 1\%$) as m_{top} is believed to be ~ 30 GeV	20%	2×10^5
1987	V_{ub}/V_{cb}	No direct prediction	0.1	2×10^5
1994	$b \rightarrow s\gamma$	$(2.8 \pm 0.8) \times 10^{-4}$	2.3×10^{-4}	4×10^6
2001	$\sin(2\beta)$	No direct prediction, but consistent with other measurements	3/4	10^7

◆ B physics is an experimentally driven field with exciting discoveries, many not predicted.

◆ There is much much more physics to do.

The Physics: General


Mysteries

Dark
Matter

Dominance
of Matter over
Antimatter


Dark
Energy

Solutions: New Physics



The Physics: More Specific

- CP Violation: Particles behave differently than antiparticles
 - Demonstrated in B decays by BaBar & Belle (one \angle measured, β)
 - But there are 4 different angles to determine: $\alpha, \beta, \gamma, \chi$
 - Different incarnations of **New Physics** affect these angles in different ways. **New Physics** can show up as inconsistencies between/among CP measurements and other quantities.
- Rare Decays




- **New Particles** can appear in the loop & interfere – Phases of the new physics can be investigated


BTeV
Co

BTeV's Staged Detector



12 9 6 3 0 3 6 9 12 m



Electromagnetic
Calorimeter



Two-component RICH



Only Top
PMTs in
Stage I

- Stage I detector
 - 50% of EM cal - *we retain 60% of the rate on neutrals*
 - No liquid radiator system - *we retain 75% of flavor tagging rate*
 - Straw stations 3 & 4 are missing, as are Silicon stations 3, 4 & 7 - *no real physics effects, these are for redundancy*
 - No dimuon trigger & only 2 muon tracking stations - *no real effects, the dimuon trigger is a useful systematic check but can come later*
 - 50% of the trigger & DAQ highways - *no real effects on b's as there is alot of "head room" in the system and we can give up some charm initially*
- Stage II detector adds in all the missing components

- Stage I starts August 1, 2009
- Then we run until July 1, 2010
 - Expect about 1 month to commission IR
 - Then its up to us to produce physics
- Summary of Stage 1
 - Estimate 6 months running time
 - Lab says that we will run 10 months a year and get 1.6 fb^{-1}
 - Thus this is a 1 fb^{-1} run
 - We have 75% of our “normal” rate on all charged flavor tagged modes
 - We have $75\% \times 60\% = 45\%$ of our “normal” rate on flavor tagged modes with neutrals
- Some Commissioning done before on wire target or at end of stores and during the 1 month IR commissioning – *New IR has $2.5 \times \mathcal{L}$ than when BTeV was approved by P5!*

LHC & LHCb's Schedule

- LHC running in steady state
 - In steady state mode, after a few years, they are scheduled to run 160 days a year for physics MINUS running for Heavy Ions - estimate 139 days on pp (see Collier, Proc. Chamonix XII, March 2003, CERN-AB-2003-008 ADM)
 - LHCb will start running at 2.8×10^{32} ; this gives using the formula in Collier 0.8 fb^{-1} per calendar year
- LHCb initial running constraints
 - Initially plan to set $\beta^* 100 \times \text{ATLAS/CMS}$, to avoid multiple interactions/crossing as 1st runs will be with 1632 ns bunch spacing to avoid necessity of crossing angle (Here LHCb needs special set up to see collisions since they are displaced by 11.2 m from interaction region center)
 - First year will see limited running at 75 ns bunch spacing; LHCb will run at $2/3 \times 10^{32}$ to avoid multiple int/xing. Second year will switch from 75 ns to 25 ns “when possible”

- LHC schedule (LHCb-1)
 - Nominal: start April 1, 2007
 - We predict LHCb 2007 integrated luminosity to be 0.1 fb^{-1}
 - Since the 1st quarter of 2008 is still in the 1st year of tuning they will collect 0.6 fb^{-1}
 - They get the full 0.8 fb^{-1} in 2009
- But - this schedule has no contingency


- Therefore we choose to set up an alternate schedule similar to the one that we have that has lots of float. A defensible schedule has ~ 12 months of float implying:
 - 0 fb^{-1} in 2007
 - 0.1 fb^{-1} in 2008
 - 0.6 fb^{-1} in 2009
 - 0.8 fb^{-1} in 2010 and beyond
- Neither for BTeV or LHCb is detector commissioning considered in what follows: we assume it will factor out of the comparisons
 - BTeV has some commissioning on wire target etc...
 - LHCb has limited accesses due to interference with ATLAS, CMS, etc..

~~BTeV~~ Co Yearly Integrated Luminosity Assumptions

fb⁻¹

	2007	2008	2009	2010	2011	2012	2013	2014	Sum
LHCb-1	0.1	0.6	0.8	0.8	0.8	0.8	0.8	0.8	5.5
LHCb-2		0.1	0.6	0.8	0.8	0.8	0.8	0.8	4.7
BTeV				1.5	1.6	1.6	1.6	1.6	7.9

~~BTeV~~ C0 Comparison I - Total number of B's to "tape"




- For BTeV we take 1/2 the nominal rate in 2010 due to the staged detector
- BTeV is better by 5x from Trigger-DAQ & 2x from running time, giving a factor of 10 $b\bar{b}$'s to tape
- e^+e^- at 1000 fb^{-1} would have $0.1 \times 10^{10} b\bar{b}$'s

Measuring γ Using $B_s \rightarrow D_s K^-$

	BTeV Stage I	BTeV Stage II	LHCb[10]
Yield (2 fb^{-1})	6,750	6,750	7,140
S/B	7	7	>1
$\epsilon \cdot D^2$	9.8%	13%	7.1%
Tagged yield (2 fb^{-1})	660	878	507
Error in γ for 2 fb^{-1}	9.4°	8.4°	14.5°
Error in γ /year (steady state)		10.9°	26.5°

From LHCb
Light TDR



~~BTeV~~ co Conclusion on Measuring γ in $B_s \rightarrow D_s K^-$

- What is a meaningful measurement of a CP violating angle?
 - Example $B^0 \rightarrow \phi K_s$ CP Asymmetry = $\sin 2\beta$
Babar: $0.47 \pm 0.34 \pm 0.07$, Belle: $-0.96 \pm 0.50 \pm 0.10$
in $J/\psi K_s$ $\sin 2\beta = 0.74 \pm 0.05$. Thus both measurements are not definitive and both have an error in $\beta \sim 14^\circ$. Need $\delta\beta < 10^\circ$ or better!
- *Thus LHCb will not likely have a meaningful measurement of γ in either of their turn on scenarios before BTeV, nor will they ever make a measurement as good as BTeV's*

Measuring α using $B^0 \rightarrow \rho\pi$


- LHCb
 - Shaslik-style Pb-scintillating fiber device, energy resolution $10\%/\sqrt{E} \oplus 1.5\%$ BTeV's is $1.7\%/\sqrt{E} \oplus 0.55\%$
 - The LHCb detector segmentation is $4 \times 4 \text{ cm}^2$ up to 90 mr, $8 \times 8 \text{ cm}^2$ to 160 mr and $16 \times 16 \text{ cm}^2$ at larger angles. (The distance to the interaction point is 12.4 m.) Thus the segmentation is comparable to BTeV only in the inner region. (BTeV has $2.8 \times 2.8 \text{ cm}^2$ crystals 7.4 m from the center of the interaction region.)
 - In 2 fb^{-1} 7260 events, $S/B < 1/7.1$, no estimate from LHCb of $\delta\alpha$, we find 11.7° from these #'s compared to BTeV Stage I 6.3°
- *Since LHCb will accumulate only half the integrated luminosity of BTeV per year, it is clear that they will not be able to make a definitive measurement of α , in fact, it is likely that they will not be able to make one at all, not surprising because of the poor energy resolution and segmentation of their calorimeter.*

Measuring χ in B_s decays

- Modes
 - BTeV uses CP eigenstates: $J/\psi \eta^{(\prime)}$
 - LHCb uses $J/\psi \phi$, VV mode so they must do a transversity analysis
- CDF & D0 get 1 $J/\psi \phi$ each per $\text{pb}^{-1} \Rightarrow \delta\chi \sim 13^\circ$ in Run II, **if** B_s mixing is also measured (sets a floor on $\int L$)

	BTeV Stage I	BTeV Stage II	LHCb[10]
Yield (2 fb^{-1})	6,800	11,340	100,000
S/B	20	20	>3
$\epsilon \cdot D^2$	9.8%	13%	5.5%
Tagged yield (2 fb^{-1})	660	1474	5500
Error in χ for 2 fb^{-1}	1.1°	0.7°	3.7°
Error in χ /year (steady state)		0.9°	5.9°

Conclusions on χ




LHCb will have a chance in 2009 of making a significant measurement of χ , if it is in excess of $\sim 20^\circ$ and they collect sufficient integrated luminosity to improve over the combined CDF & DO measurement. At the end of 2010 BTeV will have the best measurement of χ and the error will eventually be less than 0.5° .

Thus BTeV has the best chance of making a significant measurement if new physics is present and is the only detector that can measure χ if new physics doesn't make a very large contribution.

- Want to measure the polarization
- No flavor tagging here
- Define $QF = \sqrt{1000 / (\# \text{ of events})} \times \sqrt{(S + B) / S}$

	BTeV	LHCb[10]
Yield (2 fb^{-1})	2277	5546
S/B	7	>0.5
QF	0.71	0.74
Yield in 1 calendar year	1700	1660
$QF/\text{year steady state}$	0.63	1.34

- BTeV eventually overtakes LHCb



- This is LHCb's best case: They trigger on dimuons, there is no flavor tagging, and yet BTeV eventually has smaller errors

Conclusions

- The LHC turn on will be a long process by their own projections. Latest information (CMS May review), it will not start before August 2007
- LHCb will have trouble dealing with initial 75 ns running
- LHCb may get lucky and measure something “easy” like B_s mixing, if CDF & D0 don’t do it but they will have to overcome what both CDF & D0 do with B_s & what the B factories do with B^0 & B^-
- In the slightly longer term, BTeV will dominate measurements of α , γ , & χ
- After 2010 BTeV’s physics reach will dominate in all areas